

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	58.8%
	Response at Six Months	Average	32.4%
	Remission at Six Months	Average	14.7%
	Follow-up at 12 Months	Average	50.0%
	Response at 12 Months	Average	20.6%
	Remission at 12 Months	Average	5.9%
Adult Depression	PHQ-9 Utilization*	Below	57.8%
	Follow-up at Six Months	Above	81.5%
	Response at Six Months	Above	37.0%
	Remission at Six Months	Above	22.8%
	Follow-up at 12 Months	Above	78.7%
	Response at 12 Months	Above	38.9%
	Remission at 12 Months	Above	25.6%
Adolescent Mental Health and/or Depression Screening*		Above	99.1%
Colorectal Cancer Screening		Above	71.0%
Optimal Asthma Control - Adults		Above	56.4%
Optimal Asthma Control - Children		Average	55.7%
Optimal Diabetes Care		Average	47.7%
Optimal Vascular Care		Above	58.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).