

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	84.8%
	Follow-up at Six Months	Above	72.4%
	Response at Six Months	Average	22.4%
	Remission at Six Months	Average	6.9%
	Follow-up at 12 Months	Above	56.9%
	Response at 12 Months	Average	15.5%
	Remission at 12 Months	Average	12.1%
Adult Depression	PHQ-9 Utilization*	Below	56.3%
	Follow-up at Six Months	Above	82.9%
	Response at Six Months	Above	39.1%
	Remission at Six Months	Above	22.5%
	Follow-up at 12 Months	Above	77.5%
	Response at 12 Months	Above	36.5%
	Remission at 12 Months	Above	21.2%
Adolescent Mental Health and/or Depression Screening*		Above	97.4%
Colorectal Cancer Screening		Above	71.5%
Optimal Asthma Control - Adults		Above	60.0%
Optimal Asthma Control - Children		Average	62.0%
Optimal Diabetes Care		Above	48.3%
Optimal Vascular Care		Above	59.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).