

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Average	46.0%
	Response at Six Months	Average	25.0%
	Remission at Six Months	Average	15.0%
	Follow-up at 12 Months	Average	51.0%
	Response at 12 Months	Average	26.0%
	Remission at 12 Months	Average	15.0%
Adolescent Mental Health and/or Depression Screening*		Average	98.4%
Colorectal Cancer Screening		Below	62.5%
Optimal Asthma Control - Adults		Average	62.2%
Optimal Asthma Control - Children		Average	74.3%
Optimal Diabetes Care		Average	49.1%
Optimal Vascular Care		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).