

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	91.8%
	Follow-up at Six Months	Average	53.1%
	Response at Six Months	Average	16.7%
	Remission at Six Months	Average	8.3%
	Follow-up at 12 Months	Average	52.1%
	Response at 12 Months	Average	21.9%
	Remission at 12 Months	Average	13.5%
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	56.3%
	Response at Six Months	Average	34.4%
	Remission at Six Months	Above	31.3%
	Follow-up at 12 Months	Average	50.0%
	Response at 12 Months	Average	18.8%
	Remission at 12 Months	Average	9.4%
Adolescent Mental Health and/or Depression Screening*		Below	87.3%
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		Above	67.9%
Optimal Asthma Control - Children		Above	71.3%
Optimal Diabetes Care		-	-
Optimal Vascular Care		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).