

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	65.7%
	Follow-up at Six Months	Average	47.6%
	Response at Six Months	Average	15.3%
	Remission at Six Months	Average	7.3%
	Follow-up at 12 Months	Average	44.4%
	Response at 12 Months	Average	10.5%
	Remission at 12 Months	Average	4.8%
Adolescent Mental Health and/or Depression Screening*		Above	98.7%
Colorectal Cancer Screening		Average	71.0%
Optimal Asthma Control - Adults		Average	57.5%
Optimal Asthma Control - Children		Average	52.3%
Optimal Diabetes Care		Average	47.5%
Optimal Vascular Care		Average	56.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).