

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	91.7%
	Follow-up at Six Months	Average	47.0%
	Response at Six Months	Average	7.2%
	Remission at Six Months	Average	6.0%
	Follow-up at 12 Months	Above	54.2%
	Response at 12 Months	Average	14.5%
	Remission at 12 Months	Average	6.0%
Adult Depression	PHQ-9 Utilization*	Below	66.3%
	Follow-up at Six Months	Below	36.4%
	Response at Six Months	Below	13.2%
	Remission at Six Months	Below	5.3%
	Follow-up at 12 Months	Average	42.2%
	Response at 12 Months	Below	13.9%
	Remission at 12 Months	Below	6.8%
Adolescent Mental Health and/or Depression Screening*		Above	96.8%
Colorectal Cancer Screening		Below	65.4%
Optimal Asthma Control - Adults		Average	56.5%
Optimal Asthma Control - Children		Average	53.7%
Optimal Diabetes Care		Average	43.8%
Optimal Vascular Care		Average	60.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).