

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	71.4%
	Follow-up at Six Months	Average	41.7%
	Response at Six Months	Average	13.9%
	Remission at Six Months	Average	8.3%
	Follow-up at 12 Months	Average	36.1%
	Response at 12 Months	Average	16.7%
	Remission at 12 Months	Average	11.1%
Adolescent Mental Health and/or Depression Screening*		Below	80.0%
Colorectal Cancer Screening		Average	68.6%
Optimal Asthma Control - Adults		Average	50.0%
Optimal Asthma Control - Children		Average	43.3%
Optimal Diabetes Care		Average	35.8%
Optimal Vascular Care		Below	35.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).