

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	94.9%
	Follow-up at Six Months	Average	61.1%
	Response at Six Months	Average	19.4%
	Remission at Six Months	Average	11.1%
	Follow-up at 12 Months	Average	55.6%
	Response at 12 Months	Average	27.8%
	Remission at 12 Months	Average	16.7%
Adult Depression	PHQ-9 Utilization*	Average	72.1%
	Follow-up at Six Months	Average	47.0%
	Response at Six Months	Average	20.3%
	Remission at Six Months	Average	13.4%
	Follow-up at 12 Months	Average	43.6%
	Response at 12 Months	Average	15.8%
	Remission at 12 Months	Average	9.4%
Adolescent Mental Health and/or Depression Screening*		Below	67.1%
Colorectal Cancer Screening		Above	66.6%
Optimal Asthma Control - Adults		Below	31.2%
Optimal Asthma Control - Children		Average	33.3%
Optimal Diabetes Care		Below	31.8%
Optimal Vascular Care		Below	36.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).