

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	40.6%
	Response at Six Months	Average	15.6%
	Remission at Six Months	Average	3.1%
	Follow-up at 12 Months	Average	37.5%
	Response at 12 Months	Average	9.4%
	Remission at 12 Months	Average	3.1%
Adult Depression	PHQ-9 Utilization*	Below	67.8%
	Follow-up at Six Months	Below	25.5%
	Response at Six Months	Below	7.4%
	Remission at Six Months	Below	4.8%
	Follow-up at 12 Months	Below	25.5%
	Response at 12 Months	Below	10.0%
	Remission at 12 Months	Average	6.1%
Adolescent Mental Health and/or Depression Screening*		Below	86.8%
Colorectal Cancer Screening		Below	38.6%
Optimal Asthma Control - Adults		Below	19.1%
Optimal Asthma Control - Children		Below	21.0%
Optimal Diabetes Care		Average	46.6%
Optimal Vascular Care		Average	54.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).