

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	51.7%
	Follow-up at Six Months	Average	37.2%
	Response at Six Months	Average	10.3%
	Remission at Six Months	Average	3.9%
	Follow-up at 12 Months	Average	42.3%
	Response at 12 Months	Average	15.4%
	Remission at 12 Months	Average	7.7%
Adolescent Mental Health and/or Depression Screening*		Below	81.3%
Colorectal Cancer Screening		Above	68.9%
Optimal Asthma Control - Adults		Below	22.0%
Optimal Asthma Control - Children		Below	16.0%
Optimal Diabetes Care		Average	45.2%
Optimal Vascular Care		Below	41.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).