

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	80.0%
	Follow-up at Six Months	Average	45.7%
	Response at Six Months	Average	20.0%
	Remission at Six Months	Average	10.0%
	Follow-up at 12 Months	Average	38.6%
	Response at 12 Months	Average	8.6%
	Remission at 12 Months	Average	2.9%
Adult Depression	PHQ-9 Utilization*	Above	84.1%
	Follow-up at Six Months	Below	25.7%
	Response at Six Months	Below	8.8%
	Remission at Six Months	Below	4.2%
	Follow-up at 12 Months	Average	37.3%
	Response at 12 Months	Average	16.2%
	Remission at 12 Months	Average	8.5%
Adolescent Mental Health and/or Depression Screening*		Above	99.4%
Colorectal Cancer Screening		Average	66.2%
Optimal Asthma Control - Adults		Above	64.2%
Optimal Asthma Control - Children		Above	69.8%
Optimal Diabetes Care		Average	49.6%
Optimal Vascular Care		Average	65.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).