

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	80.0%
	Follow-up at Six Months	Average	60.4%
	Response at Six Months	Average	14.6%
	Remission at Six Months	Average	6.3%
	Follow-up at 12 Months	Average	50.0%
	Response at 12 Months	Average	12.5%
	Remission at 12 Months	Average	8.3%
Adult Depression	PHQ-9 Utilization*	Above	92.1%
	Follow-up at Six Months	Above	57.0%
	Response at Six Months	Above	26.4%
	Remission at Six Months	Above	15.9%
	Follow-up at 12 Months	Above	54.0%
	Response at 12 Months	Above	23.9%
	Remission at 12 Months	Average	12.7%
Adolescent Mental Health and/or Depression Screening*		Below	89.7%
Colorectal Cancer Screening		Average	68.2%
Optimal Asthma Control - Adults		Above	73.0%
Optimal Asthma Control - Children		Above	69.9%
Optimal Diabetes Care		Above	52.5%
Optimal Vascular Care		Above	63.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).