

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Average	39.1%
	Response at Six Months	Average	13.9%
	Remission at Six Months	Average	6.1%
	Follow-up at 12 Months	Below	24.4%
	Response at 12 Months	Average	7.8%
	Remission at 12 Months	Average	7.0%
Adult Depression	PHQ-9 Utilization*	Above	97.4%
	Follow-up at Six Months	Average	51.3%
	Response at Six Months	Average	20.7%
	Remission at Six Months	Average	9.4%
	Follow-up at 12 Months	Below	34.9%
	Response at 12 Months	Average	13.3%
	Remission at 12 Months	Average	6.7%
Adolescent Mental Health and/or Depression Screening*		Above	97.8%
Colorectal Cancer Screening		Above	71.4%
Optimal Asthma Control - Adults		Above	58.4%
Optimal Asthma Control - Children		Above	66.7%
Optimal Diabetes Care		Average	48.4%
Optimal Vascular Care		Average	55.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).