

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	77.1%
	Follow-up at Six Months	Average	50.0%
	Response at Six Months	Average	15.0%
	Remission at Six Months	Average	5.0%
	Follow-up at 12 Months	Below	10.0%
	Response at 12 Months	Average	5.0%
	Remission at 12 Months	Average	5.0%
Adult Depression	PHQ-9 Utilization*	Below	54.3%
	Follow-up at Six Months	Average	41.4%
	Response at Six Months	Average	14.0%
	Remission at Six Months	Average	7.0%
	Follow-up at 12 Months	Below	13.2%
	Response at 12 Months	Below	4.5%
	Remission at 12 Months	Below	2.7%
Adolescent Mental Health and/or Depression Screening*		Above	96.8%
Colorectal Cancer Screening		Above	70.6%
Optimal Asthma Control - Adults		Above	59.9%
Optimal Asthma Control - Children		Average	53.8%
Optimal Diabetes Care		Average	46.6%
Optimal Vascular Care		Average	55.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).