

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	76.0%
	Follow-up at Six Months	Below	15.4%
	Response at Six Months	Average	7.7%
	Remission at Six Months	Average	5.8%
	Follow-up at 12 Months	Average	25.0%
	Response at 12 Months	Average	9.6%
	Remission at 12 Months	Average	7.7%
Adult Depression	PHQ-9 Utilization*	Below	63.6%
	Follow-up at Six Months	Below	29.8%
	Response at Six Months	Below	8.4%
	Remission at Six Months	Below	3.9%
	Follow-up at 12 Months	Below	29.4%
	Response at 12 Months	Below	6.3%
	Remission at 12 Months	Below	3.6%
Adolescent Mental Health and/or Depression Screening*		Below	79.3%
Colorectal Cancer Screening		Above	58.2%
Optimal Asthma Control - Adults		Below	17.2%
Optimal Asthma Control - Children		Below	33.5%
Optimal Diabetes Care		Below	22.1%
Optimal Vascular Care		Below	22.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).