

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	70.0%
	Follow-up at Six Months	Average	46.0%
	Response at Six Months	Average	12.0%
	Remission at Six Months	Average	6.0%
	Follow-up at 12 Months	Average	40.0%
	Response at 12 Months	Average	16.0%
	Remission at 12 Months	Average	4.0%
Adult Depression	PHQ-9 Utilization*	Below	57.0%
	Follow-up at Six Months	Below	29.3%
	Response at Six Months	Average	10.6%
	Remission at Six Months	Average	6.6%
	Follow-up at 12 Months	Below	30.0%
	Response at 12 Months	Average	13.2%
	Remission at 12 Months	Average	8.4%
Adolescent Mental Health and/or Depression Screening*		Below	82.9%
Colorectal Cancer Screening		Above	73.2%
Optimal Asthma Control - Adults		Below	25.1%
Optimal Asthma Control - Children		Below	19.9%
Optimal Diabetes Care		Below	34.8%
Optimal Vascular Care		Below	40.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).