

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	57.5%
	Response at Six Months	Average	17.5%
	Remission at Six Months	Average	2.5%
	Follow-up at 12 Months	Average	37.5%
	Response at 12 Months	Average	15.0%
	Remission at 12 Months	Average	7.5%
Adult Depression	PHQ-9 Utilization*	Average	79.4%
	Follow-up at Six Months	Average	41.5%
	Response at Six Months	Average	16.1%
	Remission at Six Months	Average	7.6%
	Follow-up at 12 Months	Average	43.2%
	Response at 12 Months	Average	14.0%
	Remission at 12 Months	Average	7.2%
Adolescent Mental Health and/or Depression Screening*		Above	97.1%
Colorectal Cancer Screening		Below	64.0%
Optimal Asthma Control - Adults		Average	62.9%
Optimal Asthma Control - Children		Above	69.8%
Optimal Diabetes Care		Average	52.0%
Optimal Vascular Care		Average	58.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).