

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	97.2%
	Follow-up at Six Months	Average	43.9%
	Response at Six Months	Average	13.6%
	Remission at Six Months	Average	10.6%
	Follow-up at 12 Months	Average	22.7%
	Response at 12 Months	Below	3.0%
	Remission at 12 Months	Average	3.0%
Adult Depression	PHQ-9 Utilization*	Above	93.0%
	Follow-up at Six Months	Average	46.2%
	Response at Six Months	Average	18.0%
	Remission at Six Months	Average	10.8%
	Follow-up at 12 Months	Below	37.0%
	Response at 12 Months	Average	15.4%
	Remission at 12 Months	Average	11.1%
Adolescent Mental Health and/or Depression Screening*		Average	91.3%
Colorectal Cancer Screening		Above	77.1%
Optimal Asthma Control - Adults		Average	54.9%
Optimal Asthma Control - Children		Average	55.1%
Optimal Diabetes Care		Average	48.0%
Optimal Vascular Care		Average	56.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).