

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	77.3%
	Follow-up at Six Months	Average	54.9%
	Response at Six Months	Above	29.4%
	Remission at Six Months	Above	27.5%
	Follow-up at 12 Months	Average	31.4%
	Response at 12 Months	Average	9.8%
	Remission at 12 Months	Average	3.9%
Adult Depression	PHQ-9 Utilization*	Above	90.2%
	Follow-up at Six Months	Average	43.7%
	Response at Six Months	Average	20.7%
	Remission at Six Months	Average	12.9%
	Follow-up at 12 Months	Average	47.9%
	Response at 12 Months	Above	23.6%
	Remission at 12 Months	Above	16.2%
Adolescent Mental Health and/or Depression Screening*		Below	76.5%
Colorectal Cancer Screening		Below	55.2%
Optimal Asthma Control - Adults		Above	53.9%
Optimal Asthma Control - Children		Average	47.7%
Optimal Diabetes Care		Average	45.9%
Optimal Vascular Care		Average	50.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).