

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	94.2%
	Follow-up at Six Months	Average	51.4%
	Response at Six Months	Average	17.1%
	Remission at Six Months	Average	11.4%
	Follow-up at 12 Months	Average	54.3%
	Response at 12 Months	Average	21.4%
	Remission at 12 Months	Average	8.6%
Adolescent Mental Health and/or Depression Screening*		Average	85.9%
Colorectal Cancer Screening		Below	59.4%
Optimal Asthma Control - Adults		Above	67.2%
Optimal Asthma Control - Children		Average	35.0%
Optimal Diabetes Care		Average	39.1%
Optimal Vascular Care		Average	45.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).