

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Below	43.1%
	Follow-up at Six Months	Average	44.1%
	Response at Six Months	Average	15.1%
	Remission at Six Months	Average	6.5%
	Follow-up at 12 Months	Average	47.3%
	Response at 12 Months	Average	19.4%
	Remission at 12 Months	Average	10.8%
Adult Depression	PHQ-9 Utilization*	Below	66.6%
	Follow-up at Six Months	Below	31.8%
	Response at Six Months	Below	11.2%
	Remission at Six Months	Below	5.9%
	Follow-up at 12 Months	Below	27.7%
	Response at 12 Months	Below	9.1%
	Remission at 12 Months	Below	5.5%
Adolescent Mental Health and/or Depression Screening*		Below	60.4%
Colorectal Cancer Screening		Below	51.0%
Optimal Asthma Control - Adults		Below	13.8%
Optimal Asthma Control - Children		Below	13.3%
Optimal Diabetes Care		Below	37.0%
Optimal Vascular Care		Below	46.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).