

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	92.8%
	Follow-up at Six Months	Average	43.8%
	Response at Six Months	Average	14.3%
	Remission at Six Months	Average	6.7%
	Follow-up at 12 Months	Average	41.9%
	Response at 12 Months	Average	12.4%
	Remission at 12 Months	Average	4.8%
Adult Depression	PHQ-9 Utilization*	Above	80.2%
	Follow-up at Six Months	Average	49.3%
	Response at Six Months	Average	16.8%
	Remission at Six Months	Average	9.8%
	Follow-up at 12 Months	Average	48.2%
	Response at 12 Months	Average	19.4%
	Remission at 12 Months	Average	11.4%
Adolescent Mental Health and/or Depression Screening*		Average	92.4%
Colorectal Cancer Screening		Below	66.6%
Optimal Asthma Control - Adults		Below	29.0%
Optimal Asthma Control - Children		Average	44.2%
Optimal Diabetes Care		Average	42.8%
Optimal Vascular Care		Average	57.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).