

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	78.1%
	Follow-up at Six Months	Average	56.4%
	Response at Six Months	Average	10.3%
	Remission at Six Months	Average	2.6%
	Follow-up at 12 Months	Average	56.4%
	Response at 12 Months	Average	18.0%
	Remission at 12 Months	Average	10.3%
Adult Depression	PHQ-9 Utilization*	Average	75.8%
	Follow-up at Six Months	Above	62.3%
	Response at Six Months	Above	26.5%
	Remission at Six Months	Above	18.6%
	Follow-up at 12 Months	Above	57.5%
	Response at 12 Months	Above	24.8%
	Remission at 12 Months	Above	17.8%
Adolescent Mental Health and/or Depression Screening*		Average	89.4%
Colorectal Cancer Screening		Average	69.8%
Optimal Asthma Control - Adults		Below	32.6%
Optimal Asthma Control - Children		Below	30.3%
Optimal Diabetes Care		Average	46.2%
Optimal Vascular Care		Average	58.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).