

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	89.4%
	Follow-up at Six Months	Below	33.8%
	Response at Six Months	Average	9.5%
	Remission at Six Months	Below	2.5%
	Follow-up at 12 Months	Average	39.8%
	Response at 12 Months	Average	12.4%
	Remission at 12 Months	Average	9.5%
Adult Depression	PHQ-9 Utilization*	Below	72.5%
	Follow-up at Six Months	Average	50.6%
	Response at Six Months	Average	18.3%
	Remission at Six Months	Average	10.6%
	Follow-up at 12 Months	Above	52.7%
	Response at 12 Months	Average	20.3%
	Remission at 12 Months	Average	12.5%
Adolescent Mental Health and/or Depression Screening*		Below	89.6%
Colorectal Cancer Screening		Above	72.8%
Optimal Asthma Control - Adults		Below	25.2%
Optimal Asthma Control - Children		Below	22.0%
Optimal Diabetes Care		Average	47.7%
Optimal Vascular Care		Average	59.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).