

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	45.2%
	Response at Six Months	Average	16.1%
	Remission at Six Months	Average	3.2%
	Follow-up at 12 Months	Average	32.3%
	Response at 12 Months	Average	9.7%
	Remission at 12 Months	Average	3.2%
Adult Depression	PHQ-9 Utilization*	Average	79.5%
	Follow-up at Six Months	Average	45.8%
	Response at Six Months	Average	15.3%
	Remission at Six Months	Average	6.8%
	Follow-up at 12 Months	Above	56.3%
	Response at 12 Months	Average	20.3%
	Remission at 12 Months	Average	11.9%
Adolescent Mental Health and/or Depression Screening*		Average	94.3%
Colorectal Cancer Screening		Above	73.3%
Optimal Asthma Control - Adults		Average	49.7%
Optimal Asthma Control - Children		Average	44.4%
Optimal Diabetes Care		Average	47.7%
Optimal Vascular Care		Average	59.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).