

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	85.7%
	Follow-up at Six Months	Average	54.1%
	Response at Six Months	Average	15.6%
	Remission at Six Months	Average	10.1%
	Follow-up at 12 Months	Average	37.6%
	Response at 12 Months	Average	13.8%
	Remission at 12 Months	Average	7.3%
Adult Depression	PHQ-9 Utilization*	Above	84.7%
	Follow-up at Six Months	Above	54.5%
	Response at Six Months	Above	23.3%
	Remission at Six Months	Average	12.9%
	Follow-up at 12 Months	Above	55.5%
	Response at 12 Months	Above	24.2%
	Remission at 12 Months	Above	14.3%
Adolescent Mental Health and/or Depression Screening*		Below	77.4%
Colorectal Cancer Screening		Average	69.8%
Optimal Asthma Control - Adults		Below	39.9%
Optimal Asthma Control - Children		Below	37.5%
Optimal Diabetes Care		Average	46.7%
Optimal Vascular Care		Average	57.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).