

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	60.0%
	Response at Six Months	Average	23.3%
	Remission at Six Months	Average	10.0%
	Follow-up at 12 Months	Average	36.7%
	Response at 12 Months	Average	16.7%
	Remission at 12 Months	Average	10.0%
Adult Depression	PHQ-9 Utilization*	Average	72.6%
	Follow-up at Six Months	Average	54.3%
	Response at Six Months	Average	17.7%
	Remission at Six Months	Average	10.3%
	Follow-up at 12 Months	Above	58.9%
	Response at 12 Months	Average	20.0%
	Remission at 12 Months	Average	12.0%
Adolescent Mental Health and/or Depression Screening*		Above	96.7%
Colorectal Cancer Screening		Below	61.4%
Optimal Asthma Control - Adults		Average	45.5%
Optimal Asthma Control - Children		Average	34.9%
Optimal Diabetes Care		Average	45.4%
Optimal Vascular Care		Above	65.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).