

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	62.9%
	Response at Six Months	Average	25.7%
	Remission at Six Months	Average	14.3%
	Follow-up at 12 Months	Average	48.6%
	Response at 12 Months	Average	17.1%
	Remission at 12 Months	Average	14.3%
Adult Depression	PHQ-9 Utilization*	Average	75.0%
	Follow-up at Six Months	Above	60.6%
	Response at Six Months	Above	26.6%
	Remission at Six Months	Above	17.3%
	Follow-up at 12 Months	Above	60.6%
	Response at 12 Months	Above	26.6%
	Remission at 12 Months	Average	14.6%
Adolescent Mental Health and/or Depression Screening*		Above	97.4%
Colorectal Cancer Screening		Above	75.2%
Optimal Asthma Control - Adults		Average	43.7%
Optimal Asthma Control - Children		Average	32.4%
Optimal Diabetes Care		Average	50.8%
Optimal Vascular Care		Average	60.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).