

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	53.3%
	Response at Six Months	Average	10.0%
	Remission at Six Months	Average	3.3%
	Follow-up at 12 Months	Average	23.3%
	Response at 12 Months	Below	0.0%
	Remission at 12 Months	Average	0.0%
Adult Depression	PHQ-9 Utilization*	Average	77.5%
	Follow-up at Six Months	Average	41.3%
	Response at Six Months	Average	18.8%
	Remission at Six Months	Average	10.0%
	Follow-up at 12 Months	Average	36.9%
	Response at 12 Months	Below	9.4%
	Remission at 12 Months	Below	3.8%
Adolescent Mental Health and/or Depression Screening*		Average	92.0%
Colorectal Cancer Screening		Below	59.6%
Optimal Asthma Control - Adults		Below	3.5%
Optimal Asthma Control - Children		Below	2.4%
Optimal Diabetes Care		Below	33.7%
Optimal Vascular Care		Average	51.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).