

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	72.7%
	Follow-up at Six Months	Average	29.4%
	Response at Six Months	Average	10.3%
	Remission at Six Months	Average	7.4%
	Follow-up at 12 Months	Average	29.4%
	Response at 12 Months	Average	11.8%
	Remission at 12 Months	Average	7.4%
Adult Depression	PHQ-9 Utilization*	Below	60.2%
	Follow-up at Six Months	Average	44.3%
	Response at Six Months	Average	15.6%
	Remission at Six Months	Average	8.2%
	Follow-up at 12 Months	Average	48.8%
	Response at 12 Months	Average	16.4%
	Remission at 12 Months	Average	9.9%
Adolescent Mental Health and/or Depression Screening*		Above	97.0%
Colorectal Cancer Screening		Above	71.1%
Optimal Asthma Control - Adults		Above	55.1%
Optimal Asthma Control - Children		Average	48.6%
Optimal Diabetes Care		Above	47.1%
Optimal Vascular Care		Above	60.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).