

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	52.8%
	Response at Six Months	Average	11.3%
	Remission at Six Months	Average	3.8%
	Follow-up at 12 Months	Average	37.7%
	Response at 12 Months	Average	15.1%
	Remission at 12 Months	Average	9.4%
Adult Depression	PHQ-9 Utilization*	Above	89.7%
	Follow-up at Six Months	Above	58.2%
	Response at Six Months	Above	24.1%
	Remission at Six Months	Above	15.5%
	Follow-up at 12 Months	Above	55.9%
	Response at 12 Months	Average	20.8%
	Remission at 12 Months	Average	12.5%
Adolescent Mental Health and/or Depression Screening*		Above	93.8%
Colorectal Cancer Screening		Above	72.4%
Optimal Asthma Control - Adults		Above	68.6%
Optimal Asthma Control - Children		Above	69.1%
Optimal Diabetes Care		Above	52.7%
Optimal Vascular Care		Above	63.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).