

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	92.6%
	Follow-up at Six Months	Below	12.3%
	Response at Six Months	Below	3.1%
	Remission at Six Months	Average	1.5%
	Follow-up at 12 Months	Below	6.2%
	Response at 12 Months	Below	0.0%
	Remission at 12 Months	Below	0.0%
Adult Depression	PHQ-9 Utilization*	Above	92.9%
	Follow-up at Six Months	Below	30.7%
	Response at Six Months	Below	9.5%
	Remission at Six Months	Average	6.4%
	Follow-up at 12 Months	Below	26.3%
	Response at 12 Months	Below	8.8%
	Remission at 12 Months	Below	4.9%
Adolescent Mental Health and/or Depression Screening*		Above	95.5%
Colorectal Cancer Screening		Above	72.5%
Optimal Asthma Control - Adults		Below	13.5%
Optimal Asthma Control - Children		Below	24.7%
Optimal Diabetes Care		Below	40.6%
Optimal Vascular Care		Below	48.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).