

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	86.1%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	80.0%
	Follow-up at Six Months	Below	33.8%
	Response at Six Months	Below	11.3%
	Remission at Six Months	Below	5.1%
	Follow-up at 12 Months	Below	30.7%
	Response at 12 Months	Below	12.0%
	Remission at 12 Months	Average	7.9%
Adolescent Mental Health and/or Depression Screening*		Above	96.2%
Colorectal Cancer Screening		Below	61.0%
Optimal Asthma Control - Adults		Above	56.9%
Optimal Asthma Control - Children		Above	60.3%
Optimal Diabetes Care		Average	46.1%
Optimal Vascular Care		Above	60.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).