

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Average	35.3%
	Response at Six Months	Average	5.9%
	Remission at Six Months	Average	2.9%
	Follow-up at 12 Months	Average	35.3%
	Response at 12 Months	Average	11.8%
	Remission at 12 Months	Average	11.8%
Adult Depression	PHQ-9 Utilization*	Average	68.1%
	Follow-up at Six Months	Below	32.5%
	Response at Six Months	Average	13.6%
	Remission at Six Months	Average	7.8%
	Follow-up at 12 Months	Below	28.6%
	Response at 12 Months	Average	11.7%
	Remission at 12 Months	Average	6.5%
Adolescent Mental Health and/or Depression Screening*		Above	96.7%
Colorectal Cancer Screening		Average	69.1%
Optimal Asthma Control - Adults		Average	55.8%
Optimal Asthma Control - Children		Below	36.2%
Optimal Diabetes Care		Above	52.5%
Optimal Vascular Care		Average	59.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).