

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	92.3%
	Follow-up at Six Months	Average	47.6%
	Response at Six Months	Average	19.1%
	Remission at Six Months	Average	9.5%
	Follow-up at 12 Months	Average	46.4%
	Response at 12 Months	Average	19.1%
	Remission at 12 Months	Average	2.4%
Adult Depression	PHQ-9 Utilization*	Above	87.9%
	Follow-up at Six Months	Average	43.2%
	Response at Six Months	Average	14.3%
	Remission at Six Months	Average	8.2%
	Follow-up at 12 Months	Average	51.4%
	Response at 12 Months	Average	19.4%
	Remission at 12 Months	Average	10.5%
Adolescent Mental Health and/or Depression Screening*		Above	99.2%
Colorectal Cancer Screening		Average	66.4%
Optimal Asthma Control - Adults		Above	66.0%
Optimal Asthma Control - Children		Average	62.8%
Optimal Diabetes Care		Average	49.8%
Optimal Vascular Care		Average	56.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).