

Clinical Quality Measures

2022 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Average	50.0%
	Response at Six Months	Average	13.6%
	Remission at Six Months	Average	4.6%
	Follow-up at 12 Months	Average	29.6%
	Response at 12 Months	Average	6.8%
	Remission at 12 Months	Average	2.3%
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Below	36.8%
	Response at Six Months	Average	15.5%
	Remission at Six Months	Average	8.8%
	Follow-up at 12 Months	Average	36.8%
	Response at 12 Months	Average	17.6%
	Remission at 12 Months	Average	10.4%
Adolescent Mental Health and/or Depression Screening*		Average	95.2%
Colorectal Cancer Screening		Above	76.2%
Optimal Asthma Control - Adults		Below	16.4%
Optimal Asthma Control - Children		Below	19.5%
Optimal Diabetes Care		Average	44.8%
Optimal Vascular Care		Average	56.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic’s actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic’s actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic’s actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).